

Global forest land-use change from 1990 to 2010: an update to a global remote sensing survey of forests.

Authors:

D'Annunzio, Rémi

Lindquist, Erik J.

MacDicken, Kenneth G.

Introduction

Monitoring the Earth's global forest resources is important. The Global Forest Resources Assessment (FRA) team of the FAO collects and collates national statistics on forest resources on 5 - 10 year intervals to provide globally synthesized information. In FRA 2010, results from a global remote sensing survey were included as an independent means of collecting comparable time-series data on the state of the World's forests between 1990 and 2005 at the regional, climatic domain and global levels ([FAO and JRC, 2012](#)). This systematic survey of remotely-sensed data sources was updated with satellite imagery and analysis for year 2010. This note presents the latest results for the extent of forest-land and changes in forest land use for the time period 1990 to 2010. The work is the result of a partnership between FAO, its member countries and the European Commission Joint Research Centre (JRC).

Data and Methods

A systematic sampling design delivers globally consistent, statistically reliable results

Globally, over 13 500 sample sites located at the intersection of each degree of longitude and latitude (Figure 1) were analyzed. No sample sites were located at latitudes higher than 75 degrees north or south. The area surveyed provided a sampling intensity of about one percent of the global land surface. The JRC processed pan-tropical sites whereas the FAO processed sites in the boreal, temperate and sub-tropical domains.

Best-available Landsat satellite imagery was used to track change in forest area

Most of the data was provided by the USGS Landsat Global Land Survey, which is composed of the best Landsat satellite images acquired closest to the target years of 1990, 2000 and 2010. Landsat imagery was used as the primary source of data because its global coverage, long time series, spectral characteristics and 30 m spatial resolution made it suitable for the detection of tree cover and change in tree cover over time. Image samples were classified to land cover and land use using an automated, supervised approach.

Over 1 billion pixels were aggregated into polygons (minimum mapping unit of 5 hectares) and analyzed at each time interval to enable detection of forest area, forest gains and forest losses.




Figure 1. Spatial distribution of sample sites across the globe

Forest land use and land-use change reported

The forest area and forest area change results of this survey are presented as a land use, in order to be consistent with the forest definition used in FAO's Global Forest Resources Assessment country reports and national reports to the United Nations Framework Convention on Climate Change (UNFCCC).

Forest land use may include periods during which the land is devoid of tree cover, for example during cycles of forest harvesting and regeneration. In such cases, a land use is considered to be forest land use when management or natural processes will, within a reasonable time, restore tree cover to the point where it constitutes a forest.

National experts in over 100 countries reviewed data

The automated classification of the sample imagery produced spatial outputs compatible with the specific tool developed by FAO and JRC. This tool was used by over 204 national experts in 107 countries to review and revise the land cover and land use interpretation of all the sample sites.

Survey results

The area in forest land use declined between 1990 and 2010

The survey shows the total forest area in 2010 was 3.89 billion hectares, which is around 30 percent of the global land area. Between 1990 and 2010, there was a net reduction in the global forest area of around 5.3 million ha/year.

Table 1: Forest area (million ha, +/- 95% confidence interval), in 1990, 2000 and 2010, by FRA region and climatic domain. Forest area figures are presented rounded to the nearest digit with significance and thus may not be directly additive from this table.

FRA region	Climatic domain	Samples	1990	2000	2010
Africa	Subtropical	122	4 +/- 51%	5 +/- 51%	4 +/- 52%
	Tropical	2415	590 +/- 6%	580 +/- 7%	560 +/- 7%
Asia	Boreal	31	16 +/- 16%	17 +/- 15%	18 +/- 16%
	Subtropical	769	130 +/- 12%	150 +/- 11%	160 +/- 11%
	Temperate	1273	70 +/- 16%	80 +/- 15%	90 +/- 15%
	Tropical	911	310 +/- 8%	290 +/- 8%	280 +/- 9%
Europe	Boreal	294	800 +/- 5%	800 +/- 5%	790 +/- 5%
	Subtropical	94	18 +/- 26%	18 +/- 25%	18 +/- 25%
	Temperate	531	270 +/- 9%	270 +/- 9%	260 +/- 9%
North and Central America	Boreal	2777	380 +/- 2%	390 +/- 2%	380 +/- 2%
	Subtropical	368	90 +/- 13%	90 +/- 13%	90 +/- 12%
	Temperate	1593	260 +/- 6%	260 +/- 6%	250 +/- 6%
	Tropical	127	70 +/- 12%	70 +/- 12%	70 +/- 12%
Oceania	Subtropical	429	30 +/- 25%	30 +/- 25%	30 +/- 25%
	Temperate	51	21 +/- 20%	21 +/- 20%	20 +/- 20%
	Tropical	300	70 +/- 19%	70 +/- 19%	70 +/- 19%
South America	Subtropical	177	20 +/- 26%	20 +/- 25%	20 +/- 25%
	Temperate	96	13 +/- 33%	13 +/- 33%	13 +/- 33%
	Tropical	1217	820 +/- 4%	790 +/- 4%	760 +/- 4%
World	Boreal	3102	1200 +/- 3%	1200 +/- 3%	1190 +/- 3%
	Subtropical	1959	300 +/- 7%	320 +/- 7%	330 +/- 7%
	Temperate	3544	630 +/- 5%	640 +/- 5%	630 +/- 5%
	Tropical	4970	1860 +/- 3%	1790 +/- 3%	1730 +/- 3%
World		13575	4000 +/- 3%	3950 +/- 3%	3890 +/- 3%

Just over half the world's forests are in tropical or subtropical climatic domains

The world's forests are distributed unevenly with just under half the world's forests in the tropical domain (45% of total area), about one third in boreal (31%) and smaller amounts in temperate (16%) and subtropical (8%) domains.

Forest loss and gains estimated at a global scale

Worldwide, the *gross* reduction in forest land use caused by deforestation and natural disasters over the 20 year time period (15.5 million hectares per year) was partially offset by gains in forest area through afforestation and natural forest expansion (10.2 million hectares per year).





Figure 2. Forest area in 1990, 2000 and 2010, by FRA region and climatic domain

There were important regional differences in forest loss and gain

Figure 3 shows notable regional differences in the rate of change in forest area. The highest rate of forest conversion to other land uses in both periods was in South America, followed by Africa and Asia. Net forest loss in the tropical domain was steady over the study period, going from 6 million hectares per year in the 1990's to 7 million hectares per year in the 2000's.

Temperate and subtropical Asia were the only regions to show net gains in forest area.

Figure 3: Annual change in forest land-use area (1990-2010) by FRA region and climatic domain



Conclusion

The survey was designed to detect and report on forest area changes at global, regional and climatic zone scales. It complements the country-based FRA process, which collects a wide range of other information that is also needed to monitor and report on the world's forests. Importantly, the remote sensing survey provides information on changes in forest land use in different climatic domains - something which is currently impossible to obtain from a compilation of national data.

The methods developed through the remote sensing survey will be used to improve the measurement and reporting of forest area and change in forest area over time as part of the continual improvement of the FRA process and can be adapted for use at national scales. These results can be an important input to national and international reporting processes where forest area and change statistics are needed such as the Convention for Biological diversity and Reducing Emissions from Deforestation and Forest Degradation in Developing countries mechanism (REDD+) under the UN Framework Convention on Climate Change.

Detailed survey results and analyses will be incorporated into the FRA 2015 report and peer-reviewed articles.

Global Forest Resources Assessment

For more information on FRA, please go to www.fao.org/forestry/fra or contact fra@fao.org
More information on the remote sensing survey is available at
www.fao.org/forestry/fra/remotesensingsurvey/en/.